

# International GCSE in Mathematics A - Paper 2H mark scheme



Question	Working	Answer	Mark	AO	Notes
1	$2 \times 2 \times 5$ <b>or</b> $2 \times 3 \times 5$ <b>or</b> $3 \times 3 \times 5$ <b>or</b> two of 20, 40, 60 ... 30, 60, 90 ... 45, 90, 105 $2 \times 2 \times 5$ <b>and</b> $2 \times 3 \times 5$ <b>and</b> $3 \times 3 \times 5$ <b>or</b> all of 20, 40, 60, 80 ... 180 30, 60, 90 ... 180 45, 90, 105 ... 180	180	3	AO1	M1 for one of 20, 30, 45 written as product of prime factors <b>or</b> list of at least 3 multiples of any two of 20, 30, 45  M1  A1 for 180 or $2 \times 2 \times 3 \times 3 \times 5$ oe
2		$7n - 5$ oe	2	AO1	M1 for $7n + k$ ( $k$ may be zero) A1
3	$\frac{1}{2} \times (10 + 14) \times 9$ oe (= 108) '108' $\times 6$ (=648) '648' $\times 0.7$	453.6	4	AO2	M1 for area of cross section  M1 (dep on previous M1) for volume of prism M1 (independent) A1 accept 454

Question	Working	Answer	Mark	AO	Notes
4	a	$p^9$	1	AO1	B1
	b	$m^{-12}$	1	AO1	B1
	c	1	1	AO1	B1
	d	$2^{\frac{1}{3}}$	1	AO1	B1
	e	$5x + 35 = 2x - 10$ <b>or</b> $x + 7 = \frac{2x}{5} - \frac{10}{5}$ eg. $5x - 2x = -10 - 35$ <b>or</b> $7 + \frac{10}{5} = \frac{2x}{5} + x$		AO1	M1 for removing bracket or dividing all terms by 5  M1 for isolating $x$ terms in a correct equation
		-15	3		A1 dep on M1
5	14000 × 4 (=56000)			AO1	M1 NB. multiplication by 4 may occur before or after percentage decrease
	0.075 × '56000' (=4200) <b>or</b>				M1
	0.075 × 14000 (=1050)				
	'56000' - '42000' <b>or</b>				M1 (dep)
	14000 - '1050'	51 800	4		A1

Question	Working	Answer	Mark	AO	Notes
6	a	triangle with vertices (3, -1) (3, -4) (5, -4)	1	AO2	B1
	b	Rotation centre (-3, 0) 90° anticlockwise	3	AO2	B1 B1 B1 accept +90 <sup>0</sup> , 270° clockwise, −270° NB. If more than one transformation then no marks can be awarded
7	a	$4 \times 15 (=60)$ <b>or</b> $\frac{a+b+c+d}{4} = 15$ <b>or</b> $4 \times 15 - 39$	2	AO3	M1
	b	$d - a = 10$ <b>or</b> $a = 11$ <b>or</b> $a = \text{“}21\text{”} - 10$ <b>or</b> $b + c = 39 - 11 = 28$		AO3	A1 M1 ft from (a) (can be implied by 11, $b$ , $c$ , 21 <b>OR</b> $a$ , $b$ , $c$ , $d$ with $b + c = 28$ )
8		0.02 × 40 000 (=800) <b>or</b> 1.02 × 40 000 (=40800) <b>or</b> 2400  "40800"×0.02(=816) <b>and</b> "41616"×0.02(=832.32) <b>OR</b> 2448.32	3	AO1	M1
					M1 (dep) method to find interest for year 2 <b>and</b> year 3
		42448.32			A1

Question	Working	Answer	Mark	AO	Notes
9	$3x + y = 13$ <b>or</b> $6x + 2y = 26$ $- 3x - 6y = 27$ $+ x - 2y = 9$  eg. $3x - 2 = 13$ <b>or</b> $15 + y = 13$	5, -2	3	AO1	M1    multiplication of one equation with correct operation selected <b>or</b> rearrangement of one equation with substitution into second  M1    (dep) correct method to find second variable A1    for both solutions dependent on correct working
10	$\frac{14}{3} \div \frac{32}{9}$  $\frac{14}{3} \times \frac{9}{32}$ <b>or</b> $\frac{126}{27} \div \frac{96}{27}$ <b>or</b> $\frac{42}{9} \div \frac{32}{9}$	answer given	3	AO1	M1  M1  A1    correct answer from correct working
11	$(6 - 2) \times 180 (=720)$ ‘720’ – $(86 + 123 + 140 + 105)$ $(=266)$ <b>or</b> ‘720’ – 454 $(=266)$ ‘266’ $\div 2$	133	4	AO2	M1    complete method to find sum of interior angles M1    dep on 1 <sup>st</sup> method mark  M1    dep on 1 <sup>st</sup> method mark A1

Question	Working	Answer	Mark	AO	Notes
12	a	8, 25, 50, 90, 112, 120	1	AO3	B1 cao
	b		2	AO3	M1 $\pm \frac{1}{2}$ sq ft from sensible table ie clear attempt to add frequencies
				A1	ft from points if 4 or 5 correct or if all points are plotted consistently within each interval at the correct heights Accept cf graph which is not joined to the origin <b>NB</b> A bar chart, unless it has a curve going consistently through a point in each bar, scores no points.
	c		2	AO3	M1 for 60 (or 60.5) indicated on cf graph or stated A1 If M1 scored, ft from cf graph If no indication of method, ft only from correct curve & if answer is correct ( $\pm \frac{1}{2}$ sq tolerance) award M1 A1
13	$P - c = \frac{1}{2}ab^2$ $\frac{2(P - c)}{a} = b^2$	$b = \sqrt{\frac{2(P - c)}{a}}$	3	AO1	M1 Isolate term in $b$ M1 Isolate $b^2$ A1 oe with $b$ as the subject

Question	Working	Answer	Mark	AO	Notes
14	<div>a</div> <div>                     2 correct points plotted                      eg (0, 4) and (3, 0)  <math>4x + 3y = 12</math> drawn                 </div> <div>b</div> <div>  </div>		<div>2</div> <div>3</div>	<div>AO1</div> <div>AO1</div>	<div>M1</div> <div>A1</div> <div>                     B3     Correct region                       B2 for <math>x = 4</math> and <math>y = -3</math> drawn <b>and</b> consistent shading correct for at least two inequalities                       B1 for <math>x = 4</math> and <math>y = -3</math> drawn                 </div>
15	<div>a</div> <div>  </div> <div>b</div> <div> <math display="block">\frac{34}{100} \text{ oe}</math> </div> <div>c</div> <div> <math display="block">\frac{23}{46} \text{ oe}</math> </div>		<div>3</div> <div>1</div> <div>1</div>	<div>AO1</div> <div>AO3</div> <div>AO3</div>	<div>                     B3     Correct diagram                      B2 for 3 over-lapping circles with 7 in intersection <b>and</b> at least 2 other correct numbers                      B1 for 3 over-lapping circles with 7 in intersection                 </div> <div>B1     ft from diagram</div> <div>B1     ft from diagram</div>

Question	Working	Answer	Mark	AO	Notes	
16	a	$M = \frac{375}{g^3}$	3	AO1	M1	
				M1	implies first M1	
	A1		accept $M = \frac{k}{g^3}$ with $k = 375$ stated elsewhere in question			
	b		$(g = )\sqrt[3]{375 \div \left(\frac{1}{9}\right)}$ oe or $\sqrt[3]{3375}$	AO1	M1	
A1						
17	a	−3	1	AO1	B1	
	b	2	1	AO1	B1	
	c	g(2) = 6	0.75 oe	2	AO1	M1
					A1	
18	correct length scale factor	728	3	AO2	M1	
	$\text{eg } \sqrt{\frac{384}{864}}$ or $\frac{2}{3}$ or $\frac{3}{2}$				M1	for complete method
	$\left(\frac{2}{3}\right)^3 \times 2457$					

Question	Working	Answer	Mark	AO	Notes
19		E, B, D, A	3	AO1	B3 All correct B2 for 3 correct B1 for 2 correct
20	<p>a</p> $\frac{4}{9} \times \frac{3}{8}$ <p>b</p> $\frac{5}{9} \times \frac{4}{8} + \frac{4}{9} \times \frac{5}{8} \text{ or } \frac{20}{72} + \frac{20}{72} \text{ oe}$ $\text{or } 1 - \frac{4}{9} \times \frac{3}{8} - \frac{5}{9} \times \frac{4}{8} \text{ or } 1 - \frac{1}{6} - \frac{5}{9} \times \frac{4}{8} \text{ oe}$	$\frac{1}{6}$ $\frac{5}{9}$	2  3	AO3  AO3	<p>M1</p> <p>A1 oe , eg <math>\frac{12}{72}</math> Allow 0.16(666...) rounded or truncated to at least 2dp</p> <p>M2 M1 for <math>\frac{4}{9} \times \frac{5}{8}</math> or <math>\frac{5}{9} \times \frac{4}{8}</math> or <math>\frac{20}{72}</math> oe</p> <p>Accept fractions evaluated <math>\frac{20}{72} = 0.27\dot{7}</math> , <math>\frac{12}{72} = 0.16\dot{6}</math> rounded or truncated to at least 2dp</p> <p>A1 oe, e.g. <math>\frac{40}{72}</math> or <math>\frac{20}{36}</math></p>



Question	Working	Answer	Mark	AO	Notes
21	$\frac{\sin 47}{13.8} = \frac{\sin MLN}{8.5}$ $MLN = \sin^{-1}\left(\frac{\sin 47 \times 8.5}{13.8}\right)$ $MLN = 26.7(73\dots)$ $LMN = 180 - 47 - '26.7\dots' \text{ or } 106(.2260622\dots)$ $\frac{1}{2} \times 8.5 \times 13.8 \times \sin('106')$	56.3	6	AO2	<p>M1 Or method using a right angled triangle to find length <math>MX</math> (<math>MX</math> is perpendicular to <math>LN</math>)</p> $\sin 47 = \frac{MX}{8.5}$ <p>M1 Or <math>\cos^{-1} = \frac{8.5 \sin 47}{13.8}</math></p> <p>A1 <math>LMX = 63.232</math></p> <p>M1 <math>LMN = 63.232 + (180 - (90 + 47))\dots</math> or <math>106(.2260622\dots)</math></p> <p>M1</p> <p>A1 Accept an answer that rounds to 56.3 or 56.4 unless clearly obtained from incorrect working.</p>
22	<p>a</p> $2(x^2 - 4x) + 9 \text{ or }$ $2(x^2 - 4x + \frac{9}{2})$ $2((x - 2)^2 - 2^2) + 9 \text{ or }$ $2((x - 2)^2 - 2^2 + \frac{9}{2})$	$2(x - 2)^2 + 1$ <p>b explanation</p>	<p>3</p> <p>1</p>	<p>AO1</p> <p>AO1</p>	<p>M1</p> <p>M1</p> <p>A1</p> <p>B1 E.g. Because minimum is at (2, 1)</p>

Question	Working	Answer	Mark	AO	Notes
23	$\overrightarrow{BC} = \overrightarrow{BA} + \overrightarrow{AC}$ or $\begin{pmatrix} -2 \\ -3 \end{pmatrix} + \begin{pmatrix} 9 \\ 4 \end{pmatrix}$ or $\begin{pmatrix} 7 \\ 1 \end{pmatrix}$ $\sqrt{7^2 + 1^2}$	$\sqrt{50}$ oe	3	AO2	M1  M1 dep A1 accept 7.07(06...)
24	$\frac{(\sqrt{12} - 1)(2 + \sqrt{3})}{(2 - \sqrt{3})(2 + \sqrt{3})}$ $\frac{2\sqrt{12} - 2 + \sqrt{12}\sqrt{3} - \sqrt{3}}{4 - 3}$ $\sqrt{12} = 2\sqrt{3}$	shown	4	AO1	M1 method to rationalise  M1 correct expansion of brackets  B1 may be seen before expansion A1 answer from fully correct working with all steps seen
25	$(v = ) 3t^2 - 5 \times 2t - 8$ $3t^2 - 10t - 8 = 0$ $(3t + 2)(t - 4) = 0$	4	4	AO1	M1 for 2 out of 3 terms differentiated correctly A1 correct equation M1 for method to solve quadratic A1 $t = 4$ only